





DNA/RNA Blotting with the Trans-Blot® SD Semi-Dry Electrophoretic Transfer Cell



The Trans-Blot SD DNA/RNA agarose gel support frame, used with the Trans-Blot SD semi-dry electrophoretic transfer cell, allows blotting of DNA/RNA from agarose gels to nylon membrane in minutes, without any gel pretreatments. The Trans-Blot SD semi-dry electrophoretic transfer cell generates a high field strength (V/cm²) which facilitates rapid transfers. The blotting frame allows highly efficient transfers by eliminating crushing of agarose gels.

DNA Transfers in 10 Minutes

PCR* fragments, plasmid, and vector DNA ranging in size from several hundred bases to 15 kilobases can now be quickly transferred in as little as 10 minutes. Set-up time is reduced to just minutes because hydrolyzation of DNA fragments is not required prior to transfer.

RNA Transfers in 30-35 Minutes

The agarose gel support frame can also be used to transfer RNA up to 3.5 kb in 30–35 minutes, although transfer of 28s rRNA is equally efficient as determined by EtBr staining.¹

Northern and Southern Blotting

The agarose gel support frame will accommodate 6 mm thick agarose gels up to 15×20 cm. Table 1 shows the volumes of agarose required for each gel size.

Table 1. Agarose Volumes

Gel Dimensions (cm)	Volume Agarose (ml)
7 x 10	42
10 x 15	90
15 x 15	135
15 x 20	180

For convience, 10x TBE and TAE premixed buffers are avaliable. Extra thick, 2.6 mm, filter paper is necessary to provide enough buffer for transfer. The extra thick blot paper must be identical to the dimensions of the agarose gel. Bio-Rad's extra thick blot paper is available in four sizes: 7.5×10 cm, 10×15 cm, 15×15 cm, and 15×20 cm. Use of any other type of blot paper will decrease transfer efficiency.





Fig. 1. Trans-Blot SD DNA/RNA blotting kit transfer of lambda DNA fragments. A. 10 minute transfer, B. 25 minute transfer of digests of lambda DNA (NEB) and pPL-lambda (left promoter of lambda, Pharmacia). Fragment sizes were 15 kb, 4.7 kb, 1.2 kb, and 0.5 kb. Lanes contain: 1. DNA high range standard (Bio-Rad); 2. 100 ng of lambda DNA digest; 3. 50 ng; 4. 10 ng; 5. 1 ng; 6. 500 pg. DNA was electrophoresed in a 0.7% agarose TBE gel, stained, and blotted in the Trans-Blot SD semi-dry electrophoretic transfer cell. Blots were hybridized (7% SDS, 0.5 M NaPO₄, pH 8.0, 1mM EDTA) overnight at 65 °C with labeled probe (1 x 106 cpm/ml of buffer). The probe was produced by random primer labeling (Bio-Rad) of lambda PL promoter (GenBlock, Pharmacia) with 50 μCi of alpha ³²P dCTP (3,000 Ci/mM, Amersham) having a specific activity of 8.2 x 108 dpm/μg.







Fig. 2[†]. TGF-ß induced elastin expression in porcine vascular smooth muscle cells1: electrophoretic blotting using Trans-Blot SD semi-dry electrophoretic transfer cell and the Trans-Blot SD DNA/RNA blotting kit. Total RNA was extracted and processed for hybridization.¹ Each lane contained ~ 9 µg of total RNA. Confluent cultures of smooth muscle cells were treated with 5 ng/ml recombinant human transforming growth factor-B1 (TGF-B). Lanes 1 - 6 for all panels are: (1) Positive control, total RNA from neonatal porcine thoracic



aorta; (2) 0 hr after TGF-B; (3) 2 hr after TGF-B; (4) 6 hr after TGF-B; (5) 8 hr after TGF-B; (6) 12 hr after TGF-B. A. Gel after EtBr staining and 0.5x TBE equilibration. B. Membrane after 35 minute electrophoretic blotting at 3 mA/cm². The differential transfer seen between the 28s and 18s ribosomal bands can be lessened by extending the run 5-6 min. C. Autoradiograph of membrane filter hybridized with a [32P]-cDNA probe for elastin. Numbers indicate arbitrary densitometric units.

C

References

- 1. Lane, K., Zoia, O. and Davidson, J. M., Biotechniques, 12, 3, 340-346 (1992).
- 2. Trans-Blot Instruction Manual

Ordering Information		
Catalog Number	Product Description	
Support Frame		
170-4019	Agarose Gel Support Frame, 16 x 21 cm	
DNA/RNA Blotting Kit Accessories		
170-3965	Extra Thick Blot Paper, 7.5 x 10 cm, 60	
170-3958	Extra Thick Blot Paper, 10 x 15 cm, 30	
170-3959	Extra Thick Blot Paper, 15 x 15 cm, 30	
170-3960	Extra Thick Blot Paper, 15 x 20 cm, 30	
Trans-Blot SD Electrophoretic Transfer Cells		
170-3940	Trans-Blot SD Semi-Dry Electrophoretic Transfer Cell, complete unit	
170-3948	Trans-Blot SD System, 100/120 V, includes Trans-Blot SD semi-dry electrophoretic trans- fer cell and PowerPac 200 Power Supply	
170-3949	Trans-Blot SD System, 220/240 V	
Power Supplies		
165-5052	PowerPac 200 Power Supply, 100/120 V	
165-5053	PowerPac 200 Power Supply, 220/240 V	
Zeta-Probe® GT Blotting Membrane		
162-0191	Sheets , 10 x 15 cm, 15	
162-0192	Sheets , 15 x 15 cm, 15	
162-0193	Sheets, 15 x 20 cm, 15	
Premixed Buffers		
161-0733	10x Tris/Boric Acid/EDTA, 1 L	
161-0756	10x Tris/Boric Acid/EDTA, 6 x 1 L	
161-0743	50x TBS/Acetic Acid/EDTA, 1 L	
	Catalog Number Support Fra 170-4019 DNA/RNA I 170-3965 170-3958 170-3959 170-3960 Trans-Blot 170-3940 170-3948 170-3949 Power Supp 165-5052 165-5053 Zeta-Probe 162-0191 162-0192 162-0193 Premixed B 161-0733 161-0756	

For more information call your local Bio-Rad representative.

50x Tris/Acetic Acid/EDTA, 6 x 1 L

- † Used with permission of Kirk Lane, Vanderbilt University, Nashville, TN.
- Practice of PCR is covered by U.S. patent numbers 4,683,195, 4,683,202, and 4,899,818 issued to Cetus Corporation which is a subsidiary of Hoffmann-LaRoche Molecular Systems, Inc. Purchase of any of Bio-Rad's PCR-related products does not convey a license to use the PCR process covered by these patents; the user of these products to perform PCR must obtain a license.



Bio-Rad Laboratories

Life Science Group

Website www.bis-red.com Blo-Red Laboratories Main Office 2000 Affed Nobel Drive, Hercules, CA 94547, Ph. (\$70) 747-1000, Fx. (\$70747-5800) Mebelte max. bis-rad com Blo-Rad Laboratoriae Main Office 2000 Alhad Nobel Drive, Francisc, C9 94547, Fr. (5.10) 747-7004, Fr. (5.10) 747-7004, Fr. (5.10) 747-7004, Fr. (5.10) 747-7004, Fr. (6.10) 7

161-0759

BUNG 17 M US/EG PAYO 97497 1297 5/2 120797